

Estimation of sea-ice thickness and volume in the Sea of Okhotsk based on ICESat data

Sohey Nihashi¹, Nathan T. Kurtz², Thorsten Markus², Kay I. Ohshima³, Kazutaka Tateyama⁴, and Takenobu Toyota³

¹Department of Engineering for Innovation, National Institute of Technology, Tomakomai College, 443 Nishikioka, Tomakomai 059-1275, Japan

E-mail: sohey@tomakomai-ct.ac.jp

²Cryospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³Institute of Low Temperature Science, Hokkaido University, Kita-19, Nishi-8, Kita-ku, Sapporo 060-0819, Japan

⁴Department of Civil Environmental Engineering, Kitami Institute of Technology, 165 Koen-cho, Kitami 090-8507, Japan

ABSTRACT. Sea-ice thickness in the Sea of Okhotsk is estimated for 2004–2008 from ICESat derived freeboard under the assumption of hydrostatic balance. Total ice thickness including snow depth (h_{tot}) averaged over 2004–2008 is 95 cm. The interannual variability of h_{tot} is large; from 77.5 cm (2008) to 110.4 cm (2005). The mode of h_{tot} varies from 50–60 cm (2007 and 2008) to 70–80 cm (2005). Ice thickness derived from ICESat data is validated from a comparison with that observed by Electromagnetic Induction Instrument (EM) aboard the icebreaker *Soya* near Hokkaido, Japan. Annual maps of h_{tot} reveal that the spatial distribution of h_{tot} is similar every year. Ice volume of $6.3 \times 10^{11} \text{ m}^3$ is estimated from the ICESat derived h_{tot} and AMSR-E derived ice concentration. A comparison with ice area demonstrates that the ice volume cannot always be represented by the area solely, despite the fact that the area has been used as a proxy of the volume in the Sea of Okhotsk. The ice volume roughly corresponds to that of annual ice production in the major coastal polynyas estimated based on heat budget calculations. This also supports the validity of the estimation of sea-ice thickness and volume using ICESat data.

KEYWORDS: sea-ice thickness and volume, Sea of Okhotsk, ICESat

53 INTRODUCTION

54 The Sea of Okhotsk (Fig. 1) is the southernmost sea with a sizable seasonal sea-ice cover in the
55 Northern Hemisphere. Satellite observations (e.g., Gloersen and others, 1992) show that the initial freezing
56 occurs in the northern part of the sea. The ice cover maximum extent (about 1.0×10^6 km² on average) occurs
57 from the end of February to the beginning of March. The southernmost extent of sea ice reaches the coast of
58 Hokkaido around 44°N. Interannual variability of the maximum ice area is large; about 50–90% of the sea can
59 be covered with ice and the ice typically melts away by June.

60
61 Freezing, transport, and melting of sea ice cause a redistribution of heat, salt/freshwater, and
62 nutrients. In the Sea of Okhotsk, this process has been suggested to play an important role in the meridional
63 ocean circulation and accordingly the climate and biogeochemical systems of the Sea of Okhotsk and the North
64 Pacific regions. A coastal polynya formed in the northwest shelf region (Fig. 1) is the largest ice production
65 area in the Northern Hemisphere (Ohshima and others, 2016). Additionally, a large amount of brine rejection
66 associated with the active freezing forms cold and saline dense water (Shcherbina and others, 2003). This
67 dense water is thought to be a main source for ventilation of the North Pacific Intermediate Water (NPIW;
68 Talley and others, 1991; Warner and others, 1996). In this way, the sea-ice production in coastal polynyas in
69 the Sea of Okhotsk drives the overturning in the North Pacific down to intermediate depths (approximately
70 200–800 m deep). This overturning can contribute to the material cycle and the subsequent biological
71 productivity through the supply of nutrients such as iron (Nishioka and others, 2007). Recently, weakening of
72 the overturning due to global warming and associated decrease in ice production has been suggested
73 (Nakanowatari and others, 2007; Ohshima and others, 2014), thus monitoring of changes in sea-ice area and
74 volume are needed to better understand changes in the region. Sea ice formed in the coastal polynyas is
75 advected to the south by the prevailing winds and the East Sakhalin Current (Kimura and Wakatsuchi, 2000;
76 Simizu and others, 2014). This leads to a transport of negative heat and fresh water from the north to the south
77 by ice (Ohshima and others, 2003; Nihashi and others, 2012). Furthermore, sea ice also transports nutrients
78 that affects the primary production in the south (Kanna and others, 2014).

Sea-ice volume is an important parameter for understanding the redistribution processes of heat, salt/freshwater, and nutrients by sea ice (Kwok and Rothrock, 1999; Nihashi and others, 2012; Haumann and others, 2016). Knowledge of the ice volume is also crucial to evaluate the impact of global warming on sea ice. The ice volume is determined from the area and thickness. The ice area has been monitored by passive microwave satellite observations for >30 years, and a decrease trend of $-11.9 \pm 3.3\%$ decade⁻¹ was shown in the Sea of Okhotsk (Cavalieri and Parkinson, 2012). On the other hand, the ice thickness over a broad area has been hard to monitor from both in-situ and satellite observations. Thus, until now the ice volume and its trend has been unknown in the Sea of Okhotsk.

At the southernmost area along the Hokkaido coast, measurements of ice draft (ice thickness below the sea surface) had been made by moored ice-profiling sonar (IPS) observations (Fukamachi and others, 2003, 2006). The ice draft averaged over 1999–2000 is 0.6 m; the annual mean draft ranges between 0.49–0.72 m (Fukamachi and others, 2006). Similar IPS observations had been conducted near Sakhalin; the ice draft averaged over 2002–2003 is 1.05 m (Fukamachi and others, 2009). Since a coastal polynya is formed at this site, the ice draft can be separated into distinct periods of thin and thick ice. The ice draft averaged over these thin and thick periods are 0.17 m and 1.95 m, respectively. The thin ice corresponds to the coastal polynya period, while the thick ice is due to drifting of heavily deformed pack ice formed in the north. Near Hokkaido, in-situ sea-ice monitoring has been conducted every February by the icebreaker *Soya*, a patrol vessel of the Japan Coast Guard. Ice thickness was measured from the ship by video and visual observations. The video observations revealed that thickness of undeformed (level) ice averaged over 1991–2000 is 0.33 m (Toyota and others, 2004). Interannual variability of the thickness is large; annual mean thickness of level ice for the above period ranges between 0.19–0.55 m. From a comparison with the ice draft measured using moored IPS near Hokkaido (Fukamachi and others, 2003), the ice thickness measured by video observation is validated (Toyota and others, 2004). The visual observation revealed that ice thickness averaged over 2003–2005 is 111 cm (Toyota and others, 2007), with the primary difference from the video observations being due to the fact

105 that the visual observations included deformed (ridged) ice. Ice thickness measurements by Electromagnetic
106 Induction Instrument (EM) aboard the icebreaker *Soya* have been conducted since 2004 (Tateyama and others,
107 2006), and the ice thickness was validated from comparisons with that measured by the drill-hole observations
108 within the accuracy of about 10% of the thickness (Uto and others, 2006; Toyota and others, 2009).

109

110 Ice thickness estimation based on satellite observation has been also attempted in the Sea of Okhotsk.
111 Ice thickness was estimated from a radar backscatter image near Hokkaido acquired by synthetic aperture
112 radar (SAR) based on comparisons with ice thickness observed aboard the icebreaker *Soya* (Nakamura and
113 others, 2006). Toyota and others (2011) showed ice thickness distributions of the southern Sea of Okhotsk
114 during 2007–2009 based on an empirical relationship between the SAR backscatter and ice thickness (surface
115 roughness). The annual mean ice thickness was shown to range between 0.33–0.42 m. From a numerical model,
116 ice thickness distributions over the entire Sea of Okhotsk have been simulated (Watanabe and others, 2004).
117 The model results showed that the ice is thick (about 3.5 m) at the southern end of the Shantar Islands.

118

119 The Geoscience Laser Altimeter System (GLAS) instrument on the Ice, Cloud, and land Elevation
120 Satellite (ICESat) was launched on January 2003, and operated until October 2009. One of two channels of
121 GLAS (at 1064 nm) is used for surface altimetry measurement (Zwally and others, 2002). The footprint size
122 of the laser is about 70 m and the measurement interval is about 170 m along the track. Retrieval of sea-ice
123 freeboard (height of the ice plus snow surface above sea level) has been achieved from the surface elevation
124 measurements by ICESat (Kwok and others, 2004; Forsberg and Skourup, 2005; Kwok and others, 2006, 2007;
125 Markus and others, 2011).

126

127 In the Arctic Ocean, which is predominantly covered by relatively thick multi-year ice, basin-scale
128 ice thickness distribution was estimated based on ICESat derived freeboard (Kwok and others, 2009). They
129 also showed that ice drafts from ICESat and moored IPSs are consistent within 0.5 m in the Chukchi and
130 Beaufort seas. Ice thickness estimation using ICESat data in the Arctic Ocean has also been conducted by

131 Kurtz and others (2009). Zwally and others (2008) and Kern and Spreen (2015) had estimated sea-ice thickness
 132 from ICESat derived freeboard in the Weddell Sea, Southern Ocean, where relatively thick ice exists. Kurtz
 133 and Markus (2012) estimated sea-ice thickness from ICESat derived freeboard over the entire Southern Ocean,
 134 which is a typical seasonal ice zone similar to the Sea of Okhotsk (the ice thickness was found to be thinner
 135 than that in the Arctic Ocean). The objective of this study is to show the retrieval of the sea-ice thickness
 136 distribution and volume over the entire Sea of Okhotsk based on ICESat data.

137

138 METHODS AND DATA

139 A schematic illustration for the ice thickness estimation is shown in Figure 2. Total thickness of the
 140 sea ice above and below the water level (h_i) can be calculated from ICESat derived freeboard (h_f) and snow
 141 depth (h_s) under the assumption of hydrostatic balance, as follows:

142

$$143 \quad h_i = \frac{\rho_s - \rho_w}{\rho_w - \rho_i} h_s + \frac{\rho_w}{\rho_w - \rho_i} h_f. \quad (1)$$

144

145 From ice core observations (Toyota and others, 2007), snow and ice densities are assumed to be $\rho_s = 225 \text{ kg}$
 146 m^{-3} and $\rho_i = 888 \text{ kg m}^{-3}$, respectively. Sea-water density is assumed to be $\rho_w = 1026 \text{ kg m}^{-3}$ from CTD
 147 observations (Ohshima and others, 2001). Based on a relationship between ice thickness and snow depth
 148 obtained from in-situ measurements (Toyota and others, 2007), h_s is given by,

149

$$150 \quad h_s = 0.1 h_i. \quad (2)$$

151

152 The estimation of h_i highly depends on the ρ_s , ρ_i , and h_s values. Errors in the h_i estimation caused by uncertainty
 153 of these parameters are discussed in the summary and discussions section.

154

155 In this study, h_f is estimated from surface elevation measured by ICESat (h_{obs} ; Zwally and others,

2003), following a method by Kurtz and Markus (2012). Filtering of the data is first done since the h_{obs} value is significantly affected by atmospheric scattering due to cloud and blowing snow. The biased h_{obs} data is removed using instrument and waveform derived parameters that are provided along with the h_{obs} data. To obtain h_f , the local sea surface height (h_{ssh}) is subtracted from h_{obs} . The local sea surface height determined by geoid, tides, and atmospheric pressure variations (h_{est}) is first estimated, and subtracted from the h_{obs} data. By using the h_{obs} and h_{est} values, h_f is given by

$$h_f = h_{obs} - h_{est} - h_{tp}, \quad (3)$$

where h_{tp} is the sea surface tie point. The h_{tp} value is taken to be the average of the lowest three values of $h_{obs} - h_{est}$ that are within ± 12.5 km along-track spatial distance from each measurement.

RESULTS

ICESat derived h_f in the Sea of Okhotsk is shown in Figure 3 and the annual and monthly mean h_f are shown in Table 1. The data obtained from late-February to mid-March were used, except in 2007 when the data obtained from mid-March to mid-April were used. These periods are determined by the original ICESat collection periods (Zwally and others, 2003). In this study, all available spring ICESat data in the Sea of Okhotsk were used. The number of data points used indicates that the annual mean value is mainly determined by the data in March (Table 1). Sea-ice concentration derived from the Advanced Microwave Scanning Radiometer for EOS (AMSR-E) on the Aqua satellite averaged for February–March (March–April in 2007) is also shown in Figure 3. These periods correspond to the timing of the maximum ice extent. Daily mean ice concentration by Cavalieri and others (2004) were used. The ice concentration data are mapped onto a NSIDC polar stereographic grid at a spatial resolution of about 12.5 km. Sea-ice area (ice area excluding open water fraction) is also shown in Table 1. In this study, the ice edge is defined as the 30% ice concentration contour following Nihashi and others (2012). Interannual variability of the ice area is large; the largest area occurred in 2004 while the area in 2006 and 2007 was relatively small (Fig. 3). The climatology of h_f averaged over the

entire period of 2004–2008 is 18.3 cm (Table 1).

Total sea-ice thickness with snow depth ($h_{tot} = h_i + h_s$) which is calculated from Eqns (1) and (2) using h_f is also shown in Figure 3. The correspondence between h_f and h_{tot} is indicated by the color bars. The annual mean h_{tot} is thinnest in 2008 when the thickness was 77.5 cm (Table 1). The thickest ice was in 2005 when the ice thickness reached 110.4 cm. These reveal that interannual variability of ice thickness is large. No clear relationship between the sea-ice thickness and area is shown (Table 1). The climatological value of h_{tot} averaged over the entire period of 2004–2008 is 95.0 cm. The frequency histogram of h_{tot} is shown in Figure 4. The peak of the histogram (mode) varies between the 50–60 cm thickness bin (2007 and 2008) to the 70–80 cm bin (2005; Table 1).

To investigate the quality of the ice thickness retrievals from ICESat data, we compare with thickness observations by EM aboard the icebreaker *Soya* near Hokkaido. Unfortunately, the ICESat data near Hokkaido is available only three years of 2004, 2005, and 2008 (Fig. 3). The total ice thickness (h_{tot}) from EM measurements is shown in Figures. 5a, c, and e with the periods indicated in Table 2. The EM measurements have been carried out in almost the same area and period every year. Close-up maps of the ICESat derived h_{tot} corresponding to the periods of the EM measurements are shown in Figures 5b, d, and f. The periods of the ICESat measurements are also indicated in Table 2. The ICESat measurements in 2004 and 2008 were carried out about 10 days after the EM measurements while the temporal gap is 3 days in 2005. In 2004, maps of daily ice concentration (not shown) indicate that the ice edge had advanced between the EM and ICESat observations. Since the temporal gap of the measurements is short in 2005, significant differences with the sea-ice area were not seen. In 2008, the ice edge had retreated between the EM and ICESat observations.

The EM and ICESat derived h_{tot} are compared by averaging them within an analysis area indicated by a box in Figure 5, this was done because the temporal gap between EM and ICESat data is not small and their footprint size is largely different (10 m vs. 70 m). The areas close to Hokkaido, where the effects of

208 convergent ice motion and land contamination are expected to be relatively large, are excluded from the
 209 analysis area. The averaged h_{tot} with spatial standard deviation are shown in Figure 6. The interannual
 210 variability of the averaged h_{tot} derived from EM and ICESat data roughly corresponds with each other. In 2004,
 211 the averaged h_{tot} derived from EM and ICESat data are 87.5 ± 53.4 cm and 84.2 ± 35.7 cm, respectively. The
 212 mode of h_{tot} derived from the EM and ICESat data are 60–70 cm bin and 70–80 cm bin, respectively. In that
 213 year, the average and modal values are close. In 2005, the averaged h_{tot} derived from EM data is 107.6 ± 81.3
 214 cm, while that from ICESat data is 181.5 ± 69.9 cm. The mode of the ICESat derived h_{tot} is 160–170 cm, and
 215 it is close to the average value. However, the mode of the EM derived h_{tot} is 0–10 cm bin. This indicates that
 216 thin ice was dominant around the ship because the EM, whose footprint size is about 10 m, measures ice
 217 thickness at the side of the ship. On the other hand, visual observations, in which sea ice is observed in the
 218 viewing range of about 1 km, reveal that thicker ice which exceeds continuous ice breaking capability of the
 219 ship (> 1 m) was prominent. The thin modal value of h_{tot} derived from EM data (0–10 cm bin) indicates that
 220 the ship tended to look for thin ice area where the ship can break ice easily and continuously. Even so, the
 221 averaged h_{tot} in 2005 from EM data is thickest among the three years. In 2008, averaged h_{tot} derived from EM
 222 and ICESat data are 45.4 ± 39.6 cm and 70.6 ± 29.8 cm, respectively. The modal values of h_{tot} derived from
 223 EM and ICESat data are the 40–50 cm and 60–70 cm bins, respectively.

224

225 A strict comparison of h_{tot} cannot be made from Figure 6, because data of only 3 years is available
 226 and the temporal gap between EM and ICESat data reaches nearly 10 days (Table 2). In the Sea of Okhotsk,
 227 sea ice formed in the northern coastal polynyas is advected to the south by the prevailing winds and the East
 228 Sakhalin Current, and finally reaches the coast of Hokkaido (Kimura and Wakatsuchi, 2000; Simizu and others,
 229 2014). The comparison area of sea-ice thickness is surrounded by the coast of the Kuril Islands and Hokkaido
 230 (Figs. 1 and 5). Therefore, the advected sea ice is expected to drift around the analysis area for a certain period,
 231 because the sea-ice floes must pass narrow straits of the Kuril Islands to flow out into the Pacific. If the
 232 averaged h_{tot} derived from EM data represents that of the year, the relationship of h_{tot} shown in Figure 6
 233 validates the ICESat derived h_{tot} .

The ICESat derived h_{tot} (Fig. 3) is interpolated onto a polar stereographic grid at a spatial resolution of about 12.5 km using a Gaussian weighting function. This grid is same as that for the AMSR-E ice concentration data. To reveal the spatial distribution of sea-ice thickness over the entire Sea of Okhotsk for the first time, all data of each year, except for the Japan Sea, was used for the interpolation. In the Southern Ocean, a similar spatial interpolation of ICESat derived ice thickness was made by Kurtz and Markus (2012), and the influence radius was set at 125 km. In this study, the influence radius was set at 210 km because the spatial distribution of ICESat data is relatively sparse in the Sea of Okhotsk which is located at lower latitude. The spatial distribution of h_{tot} mapped onto the polar stereographic grid is shown in Figure 7. The open ocean is masked as the area where the AMSR-E ice concentration averaged from February to March (from March to April in 2007) is <30%. Ice thickness is relatively thin at the northwest shelf and north shelf regions where coastal polynyas are formed (Nihashi and others, 2009). On the other hand, ice thickness is relatively thick south of the Shantar Islands. This is considered to be caused by convergent ice motion. Similarly, ice thickness is relatively thick near Hokkaido. Ice thickness is relatively thick also in the Shelikhov and Penzhinskaya Bays. This might be caused by sea-ice deformation due to strong tides in this area (Kowalik and Polyakov, 1998). The spatial distribution of sea-ice thickness is similar every year, although the thickness is different. For example, the thickness in 2008 is thinnest over almost the entire sea-ice zone. The spatial distribution of ice thickness derived from ICESat data is roughly similar to that from a numerical model (Watanabe and others, 2004).

In 2005, when the average h_{tot} is the thickest (Table 1), the position of the Aleutian Low in January was somewhat to the south of its usual location (Figs. 8a and 8c). This indicates that the westward wind component which leads to convergent ice motion was relatively strong. The thickest ice in 2005 is considered to be caused by deformation of ice due to this anomalous wind. On the other hand, in 2008 when the average total thickness is the thinnest (Table 1), the position of the Aleutian Low in January was somewhat to the northeast of the normal location (Fig. 8b and 8c). This indicates that the eastward wind component which leads

260 to divergent ice motion was relatively strong. These results indicate that the synoptic scale wind is an important
261 factor causing interannual variability of ice thickness in the Sea of Okhotsk.

262

263 Sea-ice volume is estimated from AMSE-E derived ice area (concentration) averaged for February–
264 March (March–April in 2007) multiplied by the ICESat derived h_{tot} (Fig. 7) at each grid point. Total ice volume
265 (sum of ice volume over the sea-ice grid points) in the Sea of Okhotsk is compared with annual ice production
266 in major coastal polynyas (northwest shelf region, north shelf region, Gizhiga Bay, coastal regions of western
267 Kamchatka and northeastern Sakhalin, and Terpenia Bay; Fig. 1). This ice production is based on heat budget
268 calculation using thin ice thickness derived from AMSR-E (Nihashi and others, 2012). A time series of the ice
269 volume is shown in Figure 9. For comparison, a time series of sea-ice area (Table 1) is superimposed on the
270 figure. Variability of the ice volume roughly corresponds to that presumed from the ice area if the thickness is
271 assumed to be 1 m. However, ice volume in 2008 is relatively small owing to the thinnest ice thickness (Table
272 1). This indicates that there is a case that ice area cannot be used as a proxy of ice volume. Time series of
273 annual ice production in the major coastal polynyas is also shown in Figure 9. The ice volume and annual ice
274 production estimated from independent data are generally comparable. This also supports validity of the
275 estimations of sea-ice thickness and volume from satellite altimetry data.

276

277 SUMMARY AND DISCUSSIONS

278 In the entire sea-ice zone of the Sea of Okhotsk, the total sea-ice thickness including snow depth
279 (h_{tot}) for 2004–2008 was estimated using ICESat derived freeboard (Fig. 3), and revealed the spatial
280 distribution for the first time (Fig. 7). The h_{tot} derived from ICESat data roughly corresponded with that derived
281 from EM data near Hokkaido (Fig. 6). The errors in the h_{tot} , which is calculated from Eqns (1) and (2), are
282 mainly caused by the densities of snow (ρ_s) and sea-ice (ρ_i), snow depth (h_s), and ICESat derived freeboard
283 (h_f). In this study, densities of snow and sea-ice were assumed to be $\rho_s = 225 \text{ kg m}^{-3}$ and $\rho_i = 888 \text{ kg m}^{-3}$,
284 respectively. These values are based on in-situ measurements, and variations of ρ_s and ρ_i were estimated as \pm
285 $109 \text{ kg m}^{-3} \pm 23 \text{ kg m}^{-3}$, respectively (Toyota and others, 2007). The h_{tot} errors caused by ρ_s and ρ_i are estimated

by using these values. In the winter Arctic Ocean, relatively higher values of ρ_s (300–320 kg m⁻³) are shown (Warren and others, 1999). It is noted that the ρ_s value (225 ± 109 kg m⁻³) used in the error estimation covers these relatively higher ρ_s values in the Arctic Ocean. The ratio of h_{tot} changed by the perturbation of ρ_s and ρ_i are about $\pm 5\%$ and about $\pm 10\%$, respectively. For example, in a case of $h_{tot} = 95.0$ cm, which is estimated from a mean freeboard h_f of 18.3 cm (Table 1), the errors caused by ρ_s and ρ_i are about ± 5 cm and about ± 10 cm, respectively. In this study, snow depth h_s was assumed to be 10% of the ice thickness h_i (Eqn 2) based on in-situ measurements (Toyota and others, 2007). From the in-situ measurements of snow depth, a regression line of $h_s = 0.041 h_i + 6.070$ (cm) was shown in Toyota and others (2007). However, we didn't use this regression line, because a negative h_i value is calculated when the h_f value is small ($h_f < \text{about } 5$ cm). It is noted that h_{tot} estimated from a mean h_f of 18.3 cm using this regression line for h_s is about 93 cm, and this h_{tot} value is similar to that estimated assuming that $h_s = 0.1 h_i$ (95 cm; Table 1). When h_s is assumed that $h_s = 0.05 h_i$ based on the in-situ measurements by Toyota and others (2007), h_{tot} is estimated to be about 45% thicker than the baseline case in which h_s was assumed that $h_s = 0.1 h_i$. On the other hand, h_{tot} is estimated to be about 15% thinner than the baseline case when h_s is assumed that $h_s = 0.15 h_i$. The uncertainty of h_f has previously been shown to be 1.8 cm (Markus and others, 2011). This causes the h_{tot} error of about 9.3 cm. These estimations indicate that the h_{tot} errors owing to ρ_s , ρ_i , h_s , and h_f do not affect the validation of the ice thickness estimation shown in Figure 6.

Interannual variability of h_{tot} averaged over the entire sea-ice zone was shown to be large. The minimum value is 77.5 cm in 2008, while the maximum value is 110.4 cm in 2005 (Table 1). A frequency histogram of h_{tot} (Fig. 4) revealed that the mode varies from 50–60 cm thickness bin in 2007 and 2008 to 70–80 cm bin in 2005 (Table 1). These thickness values are much thicker than the maximum thickness of thermodynamically grown pack ice without deformation observed in the Antarctic Ocean (0.3–0.4 m; e.g. Allison and Worby, 1994; Jeffries and others, 1997; Wadhams and others, 1987). This indicates that deformed ice is prominent over the entire Okhotsk sea-ice zone as suggested from in-situ observations of sea ice in the southernmost part of the sea (Toyota and others, 2004, 2007). Furthermore, the large tails of the distributions

312 (Fig. 4) and mean thickness values of >about 80 cm (Table1) indicate that the deformed ice affects ice volume
313 in the Sea of Okhotsk.

314

315 Ice volume in the Sea of Okhotsk was estimated from ICESat derived h_{tot} and AMSR-E derived ice
316 concentration (Fig. 9). Sea-ice area has been used for a proxy of ice volume in the Sea of Okhotsk when there
317 is no ice thickness information (e.g., Nihashi and others, 2011). However, the results of this study indicated
318 that ice area cannot always represent the ice volume (Fig. 9), as in the case of 2008 when the average ice
319 thickness was thin (Table 1). The ice volume estimated in this study was shown to roughly correspond to ice
320 production in major coastal polynyas estimated based on heat flux calculations (Fig. 9). This supports the
321 validity of ice thickness and volume estimation based on ICESat data.

322

323 In the Sea of Okhotsk, sea ice formed in the northern coastal polynyas is advected to the south by
324 the prevailing northerly winds and the southward East Sakhalin Current. This indicates a transportation of
325 negative heat, fresh water, and nutrients to the south by sea ice. Simizu and others (2014) estimated the
326 southward ice-volume transport using sea-ice drift based on the moored Acoustic Doppler Current Profiler
327 (ADCP), an ocean model simulation, objective analysis data of the wind, and satellite sea-ice data. Since there
328 was no ice thickness information, they simply assumed a uniform ice thickness of 1 m. The cumulative
329 southward ice transport per winter, which crosses a line along 53°N (Fig. 1), was estimated to be 3.0×10^{11}
330 m^3 . From the climatology of ICESat derived h_{tot} of this study (Fig. 7f), h_{tot} averaged on the line is estimated to
331 be about 85 cm. Furthermore, h_{tot} averaged over the entire Sea of Okhotsk is 95 cm (Table 1). These indicates
332 that the assumption of ice thickness of 1 m in Simizu and others (2014) was almost appropriate. When the
333 ICESat derived h_{tot} (= 85 cm) is adopted to the estimation by Simizu and others (2014), the cumulative
334 southward ice transport per winter is estimated to be $2.6 \times 10^{11} \text{ m}^3$. This ice transport is comparable to the
335 climatological annual discharge of the Amur River (Fig. 1) of $3.1 \times 10^{11} \text{ m}^3$ (Fig. 1; Dai and others, 2009).

336

337 The primary goal of this study was to reveal the spatial distribution of sea-ice thickness in the Sea

338 of Okhotsk for the first time. On the other hand, in order to understand changes in sea ice associated with
339 climate change, a decadal time series of sea-ice volume is critically needed. Sea-ice area in the Sea of Okhotsk
340 has been decreasing at a rate of about 12% decade⁻¹ for the last 30 years (Cavalieri and Parkinson, 2012).
341 Furthermore, sea-ice production also has been decreasing about 11% for the last 30 years due to warming in
342 autumn at the land area northwest of the Sea of Okhotsk (Kashiwase and others, 2014). However, whether the
343 ice volume in the Sea of Okhotsk has been decreasing or not for last decades has been unknown because the
344 ICESat covers only for the period of 2004–2008. In April 2010, a radar altimeter on CryoSat-2 was launched,
345 and ice thickness and volume can be estimated also from this data (e.g., Laxon and others, 2013). By
346 combining the CryoSat-2 data (2010–) with the ICESat data, a time series of ice volume that covers >10 years
347 can be obtained. However, the CryoSat-2 data cannot be directly compared with the ICESat data because the
348 observation periods do not overlap. Furthermore, as shown in Table 2 and Figures 5 and 6, a strict comparison
349 of ice thickness between satellite and in-situ observations also cannot be made in the Sea of Okhotsk owing to
350 the large temporal gap of the measurements. Therefore, we haven't included the CryoSat-2 data in this study,
351 because reliability of the time series of ice volume requires a further in-depth study. In 2018, ICESat-2, which
352 is a successor to ICESat used in this study, is planned for launch. This data can be compared with the CryoSat-
353 2 data. In the future, analysis combining these satellite derived ice thickness would be important for discussing
354 the longer term variability of ice volume in the Sea of Okhotsk where the strong influence of global warming
355 on sea ice has been implicated.

356

357

358

359 **ACKNOWLEDGEMENTS**

360 The ICESat data were obtained from the NASA National Snow and Ice Data Center (NSIDC) Distributed
361 Active Archive Center. The AMSR-E data were provided by the NSIDC. Comments from the editor and two
362 reviewers were very helpful. This work was supported by Grant-in-Aids for Scientific Research (17K00530)
363 of the Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT).

REFERENCES

- Allison I and Worby AP (1994) Seasonal changes of sea-ice characteristics off East Antarctica. *Ann. Glaciol.*, **20**, 195–201
- Cavalieri DJ, Markus T and Comiso JC (2004) AMSR-E/Aqua Daily L3 12.5 km Sea Ice Concentration. Boulder, Colorado USA: NASA National Snow and Ice Data Center Distributed Active Archive Center (doi: 10.5067/AMSR-E/AE_SI12.003)
- Cavalieri DJ and Parkinson CL (2012) Arctic sea ice variability and trends, 1979–2010. *The Cryosphere*, **6** (4), 881–889 (doi: 10.5194/tc-6-881-2012)
- Dai A, Qian T, Trenberth KE and Milliman JD (2009) Changes in continental freshwater discharge from 1948 to 2004. *J. Clim.*, **22**, 2773–2792
- Forsberg R and Skourup H (2005) Arctic Ocean gravity, geoid and sea-ice freeboard heights from ICESat and GRACE. *Geophys. Res. Lett.*, **32**, L21502 (doi:10.1029/2005GL023711)
- Fukamachi Y, Mizuta G, Ohshima KI, Melling H, Fissel D and Wakatsuchi M (2003) Variability of sea-ice draft off Hokkaido in the Sea of Okhotsk revealed by a moored ice-profiling sonar in winter of 1999. *Geophys. Res. Lett.*, **30**(7), 1376 (doi:10.1029/2002GL016197)
- Fukamachi Y, Mizuta G, Ohshima KI, Toyota T, Kimura N and Wakatsuchi M (2006) Sea ice thickness in the southwestern Sea of Okhotsk revealed by a moored ice-profiling sonar. *J. Geophys. Res.*, **111**, C09018 (doi:10.1029/2005JC003327)
- Fukamachi Y and 8 others (2009) Direct observations of sea-ice thickness and brine rejection off Sakhalin in the Sea of Okhotsk. *Cont. Shelf Res.*, **29**, 1541–1548
- Gloersen P, Campbell WJ, Cavalieri DJ, Comiso JC, Parkinson CL and Zwally HJ (1992) Arctic and Antarctic sea ice, 1978–1987: satellite passive microwave observations and analysis. NASA SP-511, Scientific and Technical Information Program, NASA, Washington, DC
- Haumann FA, Gruber N, Münnich M, Frenger I and Kern S (2016) Sea-ice transport driving Southern Ocean salinity and its recent trends, *Nature*, **537**, 89–92 (doi:10.1038/nature19101)
- Jeffries MO, Worby AP, Morris K and Weeks WF (1997) Seasonal variations in the properties and structural

composition of sea ice and snow cover in the Bellingshausen and Amundsen Seas, Antarctic. *J. Glaciol.*,
43(143), 138–151

Kanna N, Toyota T and Nishioka J (2014) Iron and nutrient concentration in the sea ice and its impact on
surface nutritional status in the southern Okhotsk Sea. *Prog. in Ocean.*, **126**, 44–57

Kashiwase H, Ohshima KI and Nihashi S (2014) Long-term Variation in Sea Ice Production and Its Relation
to Intermediate Water in the Sea of Okhotsk. *Prog. in Ocean.*, **126**, 21–32
(doi:10.1016/j.pocean.2014.05.004)

Kern S and Spreen G (2015) Uncertainties in Antarctic sea-ice thickness retrieval from ICESat. *Annals of
Glaciology*, **56(69)**, 107–119

Kimura N and Wakatsuchi M (2000) Relationship between sea-ice motion and geostrophic wind in the
Northern Hemisphere. *Geophys. Res. Lett.*, **27(22)**, 3735–3738

Kowalik Z and Polyakov I (1998) Tides in the Sea of Okhotsk. *J. Phys. Oceanogr.*, **28**, 1389–1409

Kurtz NT, Markus T, Cavalieri DJ, Sparling LC, Krabill WB, Gasiewski AJ and Sonntag JG (2009) Estimation
of sea ice thickness distributions through the combination of snow depth and satellite laser altimetry data.
J. Geophys. Res., **114**, C10007 (doi:10.1029/2009JC005292)

Kurtz NT and Markus T (2012) Satellite observations of Antarctic sea ice thickness and volume. *J. Geophys.
Res.*, **117**, C08025 (doi:10.1029/2012JC008141)

Kwok R and Rothrock DA (1999) Variability of Fram Strait ice flux and North Atlantic Oscillation. *J. Geophys.
Res.*, **104**, 5177–5189 (doi: 10.1029/1998JC900103)

Kwok R, Zwally HJ and Yi D (2004) ICESat observations of Arctic sea ice: A first look. *Geophys. Res. Lett.*,
31, L16401 (doi:10.1029/2004GL020309)

Kwok R, Cunningham GF, Zwally HJ and Yi D (2006) ICESat over Arctic sea ice: Interpretation of altimetric
and reflectivity profiles. *J. Geophys. Res.*, **111**, C06006 (doi:10.1029/2005JC003175)

Kwok R, Cunningham GF, Zwally HJ and Yi D (2007) Ice, Cloud, and land Elevation Satellite (ICESat) over
Arctic sea ice: Retrieval of freeboard. *J. Geophys. Res.*, **112**, C12013 (doi:10.1029/2006JC003978)

Kwok R, Cunningham GF, Wensnahan M, Rigor I, Zwally HJ and Yi D (2009) Thinning and volume loss of

the Arctic Ocean sea ice cover: 2003–2008. *J. Geophys. Res.*, **114**, C07005 (doi:10.1029/2009JC005312)

Laxon SW and 14 others (2013) CryoSat-2 estimates of Arctic sea ice thickness and volume. *Geophys. Res. Lett.*, **40**, 732–737 (doi:10.1002/grl.50193)

Markus T., Massom R, Worby A, Lytle V, Kurtz N and Maksym T (2011) Freeboard, snow depth and sea-ice roughness in East Antarctica from in situ and multiple satellite data. *Annals of Glaciology*, **52(57)**, 242–248

Nakamura K., Wakabayashi H, Uto S, Naoki K, Nishio F and Uratsuka S (2006) Sea-ice thickness retrieval in the Sea of Okhotsk using dual-polarization SAR data. *Annals of Glaciology*, **44(1)**, 261–268

Nakanowatari T., Ohshima KI and Wakatsuchi M (2007) Warming and oxygen decrease of intermediate water in the northwestern North Pacific, originating from the Sea of Okhotsk, 1995–2004, *Geophys. Res. Lett.*, **34**, L04602 (doi:10.1029/2006GL028243)

Nihashi, S., Ohshima KI, Tamura T, Fukamachi Y and Saitoh S (2009) Thickness and production of sea ice in the Okhotsk Sea coastal polynyas from AMSR-E. *J. Geophys. Res.*, **114**, C10025 (doi:10.1029/2008JC005222)

Nihashi S., Ohshima KI and Nakasato H (2011) Sea-ice retreat in the Sea of Okhotsk and the ice–ocean albedo feedback effect on it. *J. Oceanogr.*, **67**, 551–562 (doi:10.1007/s10872-011-0056-x)

Nihashi S., Ohshima KI and Kimura N (2012) Creation of a heat and salt flux dataset associated with sea-ice production and melting in the Sea of Okhotsk. *J. Clim.*, **25**, 2261–2278 (doi:10.1175/JCLI-D-11-00022.1)

Nihashi S., Ohshima KI and Saitoh SI (2017) Sea-ice production in the northern Japan Sea. *Deep Sea Res. I*, **127**, 65–76 (doi:10.1016/j.dsr.2017.08.003)

Nishioka J. and 12 others (2007) Iron supply to the western subarctic Pacific: importance of iron export from the Sea of Okhotsk. *J. Geophys. Res.*, **112**, C10012 (<http://dx.doi.org/10.1029/2006JC004055>)

Ohshima KI and 7 others (2001) Winter oceanographic conditions in the southwestern part of the Okhotsk Sea and their relation to sea ice. *J. Oceanogr.*, **57**, 451–460

Ohshima KI, Watanabe T and Nihashi S (2003) Surface heat budget of the Sea of Okhotsk during 1987–2001

and the role of sea ice on it, *J. Meteorol. Soc. Jpn.*, **81**, 653–677

Ohshima KI, Nakanowatari T, Riser S, Volkov Y and Wakatsuchi M (2014) Freshening and dense shelf water reduction in the Okhotsk Sea linked with sea ice decline. *Prog. Oceanogr.*, **126**, 71–79

Ohshima KI, Nihashi S and Iwamoto K (2016) Global view of sea-ice production in polynyas and its linkage to dense/bottom water formation, *Geoscience Letters*, **3**(13) (DOI: 10.1186/s40562-016-0045-4)

Polyakov I and Martin S (2000) Interaction of the Okhotsk Sea diurnal tides with the Kashevarov Bank polynya, *J. Geophys. Res.*, **105**, 3281–3294 (doi: 10.1029/1999JC900308)

Shcherbina AY, Talley LD and Rudnick DL (2003) Direct observations of North Pacific ventilation: Brine rejection in the Okhotsk Sea, *Science*, **302**, 1952–1955

Simizu D, Ohshima KI, Ono J, Fukamachi Y and Mizuta G (2014) What drives the southward drift of sea ice in the Sea of Okhotsk? *Prog. Oceanogr.* **126**, 33–43

Talley LD (1991) An Okhotsk water anomaly: Implications for ventilation in the North Pacific, *Deep Sea Res.*, **38**, suppl. 1, S171–S190

Toyota T, Kawamura T, Ohshima KI, Shimoda H and Wakatsuchi M (2004) Thickness distribution, texture and stratigraphy, and a simple probabilistic model for dynamical thickening of sea ice in the southern Sea of Okhotsk. *J. Geophys. Res.*, **109**, C6 (doi: 10.1029/2003JC002090)

Toyota T, Takatsuji S, Tateyama K, Naoki K and Ohshima KI (2007) Properties of Sea Ice and Overlying Snow in the Southern Sea of Okhotsk. *J. Oceanogr.*, **63**, 393–411

Toyota, T., K. Nakamura, S. Uto, K.I. Ohshima, and N. Ebuchi (2009) Retrieval of sea ice thickness distribution in the seasonal ice zone from air-borne L-band SAR. *Int. J. Remote Sensing*, **30**(12), 3171–3189

Toyota T, Ono S, Cho K and Ohshima KI (2011) Retrieval of sea-ice thickness distribution in the Sea of Okhotsk from ALOS/PALSAR backscatter data. *Annals of Glaciology*, **52**(57), 177–184

Tateyama K, Shirasawa K, Uto S, Kawamura T, Toyota T and Enomoto H (2006) Standardization of electromagnetic-induction measurements of sea-ice thickness in polar and subpolar seas. *Annals of Glaciology*, **44**(1), 240–246

- 468 Uto S, Toyota T, Shimoda H, Tateyama K and Shirasawa K (2006) Ship-borne electromagnetic induction
469 sounding of sea-ice thickness in the southern Sea of Okhotsk. *Annals of Glaciology*, **44(1)**, 253–260
- 470 Wadhams P, Lange MA and Ackley SF (1987) The ice thickness distribution across the Atlantic sector of the
471 Antarctic Ocean in midwinter. *J. Geophys. Res.*, **92(C13)**, 14,535–15,552
- 472 Warner MJ, Bullister JL, Wisegarver DP and Gammon RH (1996) Basin-wide distributions of
473 chlorofluorocarbons CFC-11 and CFC-12 in the North Pacific: 1985–1989, *J. Geophys. Res.*, **101**,
474 20,525–20,542
- 475 Warren SG, Rigor IG, Untersteiner N, Radionov V.F, Bryazgin NN, Aleksandrov YI and Colony R (1999),
476 Snow depth on Arctic sea ice. *J. Clim.*, **12**, 1814–1829
- 477 Watanabe T, Ikeda M and Wakatsuchi M (2004) Thermohaline effects of the seasonal sea ice cover in the Sea
478 of Okhotsk. *J. Geophys. Res.*, **109**, C09S02 (doi:10.1029/2003JC001905)
- 479 Zwally HJ and 15 others (2002) ICESat’s laser measurements of polar ice, atmosphere, ocean, and land. *J.*
480 *Geodyn.*, **34**, 405–445 (doi:10.1016/S0264-3707(02)00042-X)
- 481 Zwally HJ and 7 others (2003) GLAS/ICESat L2 Sea Ice Altimetry Data V531, Oct. 2003 to March 2008,
482 <http://nsidc.org/data/gla13.html>, Natl. Snow and Ice Data Cent., Boulder, Colo.
- 483 Zwally HJ, Yi D, Kwok R and Zhao Y (2008) ICESat measurements of sea ice freeboard and estimates of sea
484 ice thickness in the Weddell Sea. *J. Geophys. Res.*, **113**, C02S15 (doi:10.1029/2007JC004284)

494
495
496
497
498
499
500
501

Table 1 Summary of statistics of sea-ice parameters. Here N is number of ICESat data points. The terms h_f and h_{tot} indicate freeboard and total ice thickness derived from ICESat data, respectively. For h_f and h_{tot} , those averaged over the entire sea-ice zone with spatial standard deviation are shown. For h_{tot} , the modal values are also shown. The term S_i indicates net sea-ice area which is given by excluding the open water fraction. For calculating S_i , ice concentration derived from AMSR-E data is used. The term V_i indicates sea-ice volume obtained from the ICESat h_{tot} and AMSR-E ice concentration. The term Clim. in the bottom row indicates climatology averaged for 5-years of 2004–2008 with annual standard deviation.

		ICESat	ICESat	ICESat		AMSR-E	ICESat/AMSR-E
		N	h_f	h_{tot}		S_i	V_i
Year	Month		Ave. (cm)	Ave. (cm)	Mode (cm - bin)	($\times 10^{11}$ m ²)	($\times 10^{11}$ m ³)
2004	Feb–Mar	64448	18.0 ± 12.6	93.3 ± 65.0	60–70	7.8	7.4
	Feb	15830	18.7 ± 13.3	96.9 ± 68.9	50–60		
	Mar	48618	17.8 ± 12.3	92.1 ± 63.6	60–70		
2005	Feb–Mar	68525	21.3 ± 15.4	110.4 ± 79.4	70–80	6.4	7.1
	Feb	21426	19.5 ± 13.7	100.9 ± 70.8	60–70		
	Mar	47099	22.2 ± 16.0	114.7 ± 82.7	70–80		
2006	Feb–Mar	40311	18.1 ± 13.2	93.8 ± 68.1	60–70	5.9	5.6
	Feb	9511	19.2 ± 14.7	99.3 ± 76.0	60–70		
	Mar	30800	17.8 ± 12.6	92.1 ± 65.3	60–70		
2007	Mar–Apr	36018	19.3 ± 13.1	99.9 ± 67.8	50–60	5.2	5.5
	Mar	28535	18.8 ± 13.1	97.3 ± 67.6	50–60		
	Apr	7483	21.2 ± 13.0	109.9 ± 67.4	70–80 and 80–90		
2008	Feb–Mar	37235	15.0 ± 10.0	77.5 ± 51.6	50–60	7.4	5.8
	Feb	17036	15.6 ± 10.1	80.7 ± 52.4	60–70		
	Mar	20199	14.5 ± 9.8	74.9 ± 50.7	50–60		
Clim.			18.3 ± 2.3	95.0 ± 12.0		6.5 ± 1.1	6.3 ± 0.9

502
503
504
505
506
507
508
509
510
511

Table 2 The periods of EM observations aboard the icebreaker “Soya” near Hokkaido

date of ICESat measurements in the area.		
	EM	ICESat
2004	8–12 Feb.	20 Feb.
2005	14–16 Feb.	18 Feb
2008	10–11 Feb.	20 Feb.

512




Fig. 1 Map of the Sea of Okhotsk with bottom topography. The 200- and 3000-m isobars are indicated by thin and thick solid lines, respectively. A box denotes the enlarged portion in Figure 5. White shading indicates sea-ice area (ice concentration $\geq 30\%$) in February averaged for 2003–2011; blue shading indicates open ocean area. Ice concentration from AMSR-E is used. Color shadings indicate cumulative ice production in coastal polynyas during winter (December–March) averaged from the 2002/03 to 2009/10 seasons (modified from Nihashi and others 2012; 2017). The amount is indicated by the bar scale.




Fig. 2 Schematics of sea-ice for estimation of total thickness of ice above and below the water level (h_i) from ICESat derived freeboard (h_f). Here, h_s is snow depth and h_{tot} ($=h_s + h_i$) is total ice thickness including snow depth.




Fig. 3 ICESat derived freeboard (h_f) and total ice thickness (h_{toi} ; color) superimposed on AMSR-E derived ice concentration (gray shadings) averaged for February–March (March–April in 2007), when the ICESat data was acquired. In 2004, 2007, and 2008, the Kashevarov Bank polynya, which is a sensible heat polynya (Polyakov and Martin, 2000), formed at the Kashevarov Bank region (Fig. 1).




Fig. 4 Frequency histograms of total ice thickness (h_{tot}) derived from ICESat. The histogram bin size is 10 cm. n indicates the total number of data points.




Fig. 5 **a, c, and e** Total ice thickness (h_{tot}) derived from EM measurements onboard the icebreaker “Soya”. **b, d, and f** Freeboard h_f and h_{tot} derived from ICESat measurements. The observation periods are indicated in Table 2. Sea-ice concentration derived from AMSR-E of the corresponding period is indicated by gray shadings. A box denotes the analysis area for Figure 6.




Fig. 6 A scatter plot of total ice thickness (h_{tot}) derived from EM and ICESat data. The ice thickness was averaged over the analysis area shown in Figure 5. Error bars indicate the spatial standard deviations of h_{tot} .




Fig. 7 ICESat derived total ice thickness h_{tot} interpolated onto the polar stereographic grid at the 12.5 km spatial resolution. **f** Climatology of h_{tot} averaged for 2004–2008. A black line denotes an analysis area along 53°N used in Summary and Discussions. The sea-ice data in the Japan Sea is masked out.




Fig. 8 **a** Sea level pressure (solid lines), geostrophic wind (vectors), and air temperature at 2 m (colors), averaged for the sea-ice advance season (January) in 2005. **b** As in **a**, but for data in 2008. **c** As in **a**, but for data averaged for 2004–2008. We used near-surface atmospheric data from the 6-hourly ECMWF interim reanalysis (ERA-Interim) dataset with a spatial resolution of $0.5^\circ \times 0.5^\circ$.




Fig. 9 Time series of ice volume (black solid line with triangles), ice area (gray solid line with dots), and ice production in major coastal polynyas (gray dashed line with squares) in the Sea of Okhotsk.